
Twenty First Annual University of North Georgia Mathematics Tournament

You may write in this test booklet. Only the electronic form will be graded. Correct answers are awarded one point. Incorrect or blank answers are awarded 0 points.



- a) 32
- b) $\frac{1}{4}$
- c) 25
- d) 16
- e) None of the above
- 2. A sector of a circle has angle à Find the value of àáin radians, for which the ratio of the
 - a) 2.3
 - b) 2
 - c)

- 3. Find the definite integral: $\int_{2}^{1} \frac{1}{1} \frac{x^{2}}{2^{x}} dx$
 - a) $3 2^2$
 - b) 3 2²
 - c) —
 - d) $\frac{28}{3}$
 - e) None of the above
- 4. Find the limit: $\lim_{x} x \quad x^2 \ln \frac{1 \quad x}{x}$
 - a) $\frac{1}{3}$
 - b) 0
 - c) $\frac{1}{2}$
 - d) $\frac{2}{3}$
 - e) None of the above

- 5. A Norman window is constructed by adjoining a semicircle to the top of an ordinary rectangular window. Find the dimensions (length and width) of the rectangular portion of a Norman window of maximum area if the total perimeter is 26 feet.
 - a) length $\frac{26}{4}$, width $\frac{26}{4}$
 - b) length $\frac{52}{4}$, width $\frac{26}{4}$
 - c) length $\frac{52}{8}$, width $\frac{52}{8}$
 - d) length $\frac{52}{8}$, width $\frac{26}{8}$
 - e) None of the above
- 6. Find the definite integral: $\int_{0}^{\frac{\pi}{2}} \frac{dx}{\sqrt{\sin x} \sqrt{\cos x}^{4}}$.
 - a) $\frac{1}{7}$
 - b) $\frac{1}{7}$
 - c) $\frac{1}{3}$
 - d) $\frac{\sqrt{2}}{5}$
 - e) None of the above
- 7. The area between the graph of $y ext{ ln } 1/x$, the x-axis, and the y-axis is revolved around the x-axis. Find the volume of the solid it generates.
 - a) $\frac{}{3}$
 - b)
 - c) $\frac{2}{3}$
 - d) 2
 - e) None of the above

- 8. Find the limit: $\lim_{n} \frac{n}{n^2} \frac{n}{n^2 \cdot 1} \frac{n}{n^2 \cdot 4} \cdots \frac{n}{n^2 \cdot (n \cdot 1)^2}$
 - a) $\frac{}{4}$
 - b) $\frac{}{2}$
 - c) $\frac{3}{4}$
 - d)
 - e) None of the above
- 9. Find the definite integral: $\int_{0}^{4} \sec^{3} x \, dx$
 - a) $0.5 \sqrt{2} \ln 2$
 - b)
 - c) $0.5 \sqrt{2} \ln 3$
 - d) $0.5 \sqrt{2} \ln \sqrt{2} 1$
 - e) None of the above
- 10. A person 2 meters tall walks directly away from a streetlight that is 8 meters above the rate (in meters per second)

- 11. The curve x^4y 12y 144 has one maximum point and two points of inflection. Find the area of the triangle formed by the tangents to the curve at these three points. Answers are reported in square units.
 - a) 1
 - b) 2
 - c) $\frac{4}{3}$

 - e) None of the above
- 12. Find the limit: $\lim_{x} \frac{\ln 1}{\ln 1} \frac{3^{x}}{2^{x}}$

 - b) $\frac{3}{2}$
 - c) 1
 - ln 3
 - e) None of the above
- 13. Find the integral:
 - a) $x \ln x$ $x \frac{1}{2} x \ln x$ Cb) $\ln \frac{1}{2} \frac{2x}{2}$

 - c) $x \ln x + x + \frac{1}{2} e^x + e^{2x} + x^x + C$
 - d) $x \ln x$ $x + \frac{1}{2} x \ln x$ $x^x + C$
 - e) None of the above

14. Find the integral:
$$\frac{\cos^2 x}{1 + \sin^2 x} dx$$

a)
$$\sqrt{2} \arctan \sqrt{2} \tan x \quad x \quad C$$

b)
$$2\arctan 4\tan x x C$$

c)
$$\sqrt{2} \arctan 2 \tan x \quad x \quad C$$

d)
$$4\arctan 2\tan x \quad x \quad C$$

15. Find the minimum value of the function $y = 2x^2 + 34x + 34 + e^{x-34}$.

a)
$$\frac{18}{e^{34}}$$

b)
$$\frac{26}{e^{36}}$$

c)
$$\frac{18}{e^{28}}$$

d)
$$\frac{34}{e^{34}}$$

16. Order the following six functions from slowest growing to fastest growing as x

(i)
$$2^{5x}$$
 (ii) x^x (iii) $\ln(7x)^{-7x}$ (iv) $1 e^{x/3}$ (v) $\sqrt{x^2 + 5}$

17. Evaluate the integral in terms of natural logs.

$$\int_{1/13}^{3/17} \frac{dx}{x\sqrt{1 + 25x^2}}$$

- a) ln 3
- b) $\frac{1}{5} \ln \frac{3}{17} + \frac{1}{5} \ln \frac{1}{13}$
- c) $\frac{1}{5}$ ln 13
- d) $\frac{1}{5} \ln \frac{3}{17} \ln \frac{1}{13}$
- e) None of the above
- 18. A worker ant at a super colony makes a long walk to bring food to her colony. She starts out at her mount located at the point $A \ln 2, 2\sqrt{2}$ along the curve $r_1(\) \ \sqrt{2}e$,
 - 0 $\ln 2$. At the end of this path, she then walks the straight path r_2 that connects the point B $0,\sqrt{2}$ down to the origin O(0,0). This leads to the final trail $r_3(\)$
 - $0 \sqrt{21}$. Find the total distance walked by this ant if is in radians.
 - a) $3 \ 5\sqrt{2}$
 - b) \sqrt{\sqrt{}}
 - c) 41 $\sqrt{2}$
 - d) 39 $2\sqrt{2}$
 - e) None of the above
- 19. Calculate the n^{th} derivative for f(x) $x^n e^x$ at x = 0.
 - a) ———
 - b) 0
 - c) $n(n \ 1)(n \ 2) \cdots 3 \ 2 \ 1$
 - d) $n \ln(n e)$
 - e) None of the above

20. Let f be a real-valued function defined on the interval 1,1 such that

for all x in 1,1 and let f^{-1} be the inverse function of f.

Then f^{-1} (2)

- a) 1
- b) $\frac{1}{3}$
- c) $\frac{1}{e}$
- d) $\frac{1}{2}$
- e) None of the above
- 21. Suppose f(1) = 1, f(1) = 2, g(1) = 3, and g(1) = 4. Let $h(x) = \frac{f(x)}{g(x)}$. Find h(1).
 - a) $\frac{2}{9}$
 - b) $\frac{2}{9}$
 - c) $\frac{1}{9}$
 - d)
 - e) None of the above

22. Find the definite integral:
$$\int_{0}^{1} x^{3m} x^{2m} x^{m} 2x^{2m} 3x^{m} 6^{\frac{m}{m}} dx$$

a)
$$\frac{(11)^{\frac{m-1}{m}}}{33(m-1)}$$

b)
$$\frac{(6)^{\frac{m-1}{m}}}{11(m-1)}$$

c)
$$\frac{(33)^{\frac{m-1}{m}}}{11(m-1)}$$

$$d) \quad \frac{(11)^{\frac{m-1}{m}}}{6(m-1)}$$

e) None of the above

- a) 0
- b) -
- c) $\frac{1}{3}$
- d) Does not exist
- e) None of the above

$$\int_{1}^{x^2} \ln t \ dt$$

- 24. Find the limit: $\lim_{x \to 0} \frac{1}{x^3}$.
 - a)
 - b)
 - c) 0
 - d) 2
 - e) None of the above
- 25. Suppose f is a continuous function and f(x) exists everywhere. If f(2) 10 and f(x) 3 for all x, then what is the smallest possible value for f(4)?
 - a) 1
 - b) 2
 - c) 3
 - d) 4
 - e) None of the above
- 26. Which of the following statements are true?
 - I. If f is differentiable at a, then it has a limit at a.
 - II. $\frac{x^2 + 4}{x + 2} = x + 2$
 - III. \int_{25x} is differentiable everywhere.
 - IV. _____
 - a) I and II
 - b) II and III
 - c) I and III
 - d) I and IV
 - e) None of the above

Reminder

Question 30 will be used as a tie-breaker, if necessary.

30. Write a formula for the second derivative of the composition $f \circ g$ (x) using f, g, f, g, f, g.

- a) f(x)g(x)
- b) $f g(x) g(x)^{2} f g(x) g(x)$
- c) $f g(x) g(x)^2 f g(x) g(x)$
- d) f g(x) g(x) f g(x) g(x)
- e) None of the above

31. Find the definite integral: $\int_{\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{x^2 \sin x}{1 + 2 \cos^2 x} dx$

- a) $\frac{\sqrt{3}}{9}$
- b) $\frac{\sqrt{3}}{3}$
- c) 0
- d) $\frac{5}{3}$
- e) None of the above

32. Find the smallest value of the constant k such that f(x) kx 1 $\frac{1}{x}$ 0 for all x 0.

- a) 4
- b) $\frac{1}{4}$
- c) 16
- d) $\frac{1}{16}$
- e) None of the above

33. Find the limit: $\lim_{x} \sqrt{x^{200} + x^{100} + 1} = x^{100}$

- a) 3
- b) $\frac{1}{2}$
- c) 6
- d) $\frac{1}{6}$
- e) None of the above

34. Find the definite integral: $\int_{0}^{1} \frac{x^4(1-x)^4}{1-x^2} dx$.

- a)
- b) 7
- c) 7
- d) $\frac{22}{7}$
- e) None of the above


35. Let f(x) $\frac{\cos x}{-x} \quad \text{if } x \quad \frac{\pi}{2}$ $a^2 \quad a \quad 1 \quad \text{if } x \quad \frac{\pi}{2}$

For which value of a is the function f continuous at $\frac{1}{2}$?


- a) *a* 1
- b) a 2
- c) a = 0 and a = 1
- d) a 2
- e) None of the above

36. Identify the graph of k(x) $x^{2/3}$ x^2 4.


a)




b)



c)



d)



e) None of the above

37. Find the maximum value of the function $f(x) = \frac{\sin x}{\sin^2 x} \frac{1}{\sin x}$.

- a) The maximum value of f is 0.
- b) The maximum value of f is 1.
- c) The maximum value of f is 4.
- d) The maximum value of f does not exist.
- e) None of the above

38. A cylindrical tank 10 feet high with radius 6 feet is full of water. Set up an integral to find the work required to empty the tank. The water weight is 62.4 lb/ft³.

- a) $\int_{0}^{10} 62.4$ $6^2 x dx$, where x is the distance to the top
- b) $\int_{0}^{10} 62.4$ $6^{2}(10 x) dx$, where x is the distance to the bottom
- c) Both a) and b) are correct
- d) Only a) is correct and b) is incorrect
- e) None of the above

39. Find the area enclosed between the line y = 2x and the parabola $y = x^2$.

- a) $\frac{3}{4}$
- b) $\frac{4}{5}$
- c) $\frac{4}{3}$
- d) $\frac{5}{3}$
- e) None of the above

40. If $A = \int_0^{2} \frac{\sin^n x}{\sin^n x - \cos^n x} dx$ for any n = 0, then

- a) $A = \frac{1}{2}$
- b) $A = \frac{1}{4}$